



Compendium of Veterinary Products - US edition

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Nobivac® Feline 1-HCP

Intervet/Merck Animal Health

FELINE RHINOTRACHEITIS-CALICI-PANLEUKOPENIA VACCINE

Modified Live Virus

FOR ANIMAL USE ONLY

IN THE ABSENCE OF A VETERINARIAN-CLIENT-PATIENT RELATIONSHIP, FEDERAL REGULATIONS PROHIBIT THE RELABELING, REPACKAGING, RESALE OR REDISTRIBUTION OF THE INDIVIDUAL CONTENTS OF THIS PACKAGE.

THIMEROSAL FREE

PRODUCT DESCRIPTION:

Nobivac® Feline 1-HCP vaccine is a modified live virus vaccine for the vaccination of healthy cats 9 weeks of age or older as an aid in the prevention of disease caused by feline rhinotracheitis, calici and panleukopenia viruses.

DOSAGE AND ADMINISTRATION:

Transfer the contents of the sterile diluent vial to the Nobivac Feline 1-HCP vial aseptically. Mix gently until dissolved. Use entire contents immediately after rehydration.

Inject one dose (1 mL) subcutaneously or intramuscularly at 9 weeks of age or older and repeat with a second dose 3 to 4 weeks later.

Annual revaccination with one dose is recommended. Following the advice of the American Veterinary Medical Association on Vaccination Principles, an appropriate revaccination program for individual animals should be made based upon veterinarian-client-patient relationships.

PRECAUTIONS:

- Store at 2° to 7°C (35° to 45°F).
- Do not freeze.
- Do not mix with other vaccines.
- Use new, non-chemically sterilized needles and syringes.
- Burn vaccine container and all unused contents.

WARNINGS:

- Contains gentamicin as a preservative.
- Some reports suggest that in cats, the administration of certain veterinary biologicals may induce the development of injection-site fibrosarcomas.
- Do not vaccinate pregnant cats.
- The use of a biological product may produce anaphylaxis and/or other inflammatory immune-mediated hypersensitivity reactions. **Antidote:** Epinephrine, corticosteroids, and antihistamines may all be indicated depending on the nature and severity of the reaction.

Intervet Inc., Merck Animal Health, division of Intervet Inc., Omaha, NE 68103 USA

U.S. Veterinary License No. 165A

1 800 224-5318 (USA)

1 866 683-7838 (Canada)

For patent information:

<http://www.merck.com/product/patent/home.html>

Code

25x1 DOSE 065264 141890-04

CPN: 1047049.7

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Updated March 23, 2018

Copyright 2018 - Animalytix LLC